

Working Memory Relationships with Cognitive Flexibility and Planning: A Neurocognitive Perspective

<https://doi.org/10.57642/AJOPSY1012>

Safae Sedjari

safae.sedjari@usmba.ac.ma

Department of psychology, Faculty of Letters and Human Sciences Dhar El Mahraz, Sidi Mohamed Ben Abdellah University, Fes, Morocco.

Reçu : 11/02/2025

Mohammed El-Mir

mohammed.elmir1@usmba.ac.ma

Abdellah University, Fes, Morocco.

Accepté : 11/04/2025

Publié : 15/04/2025

Abstract

The relationship between working memory and executive functions is widely recognized. Some authors describe working memory as the basis of cognitive functioning, suggesting that its capacity is a strong predictor of higher-level cognition. This relationship has also been explored at the neurological level in several studies using imaging techniques. However, this article will particularly focus on the neurocognitive relationship between working memory and two key executive functions: cognitive flexibility and planning. The aim is to highlight how working memory capacity influences cognitive flexibility and planning, and to emphasize that these three functions share common neural networks.

Keywords: working memory, cognitive flexibility, planning, neurocognitive perspective

Introduction

The relationship between performance in working memory (WM) tasks and high-level cognitive processes has been highlighted in a significant amount of research (e.g., Conway et al., 2005; Ger & Roebers, 2023; Redick et al., 2012; Unsworth et al., 2005). There is a close link between executive functions (EFs) and WM. The unitary model that emerged at the beginning of research on EFs proposed grouping all executive processes into a single component. This is evident in Baddeley's (1986) model, where the functions of the central executive play the same role as EFs. Similarly, in Cowan's (1995) model, attentional focus performed an executive role in managing attentional resources. Furthermore, the unitary model of Norman and Shallice (1986) proposed that the Supervisory Attentional System governs the entire executive operation. Miyake et al. (2000), with their integrative approach, also proposed three distinct but interconnected EFs, operating through the active maintenance of goals and relevant information during executive control. Some authors describe WM as the basis of cognitive control (Kimberg et al., 1997). It has been shown that WM capacity is correlated with performance on cognitive control tasks (e.g., Conway et al., 2001; McCabe et al., 2010; Redick et al., 2011; Schelble et al., 2012; Unsworth et al., 2012; Wilhelm et al., 2013). Ackerman et al. (2005) suggest that WM capacity is a strong predictor of high-level cognition, despite the lack of consensus about the nature of the link between these two constructs. Studies have also shown that high WM capacity is correlated with effective cognitive control, including the ability to maintain and execute goals for task completion (e.g., McVay & Kane, 2012a). In the same context, a series of studies examining the relationship between WM capacity and stimulus response time found that low WM capacity is associated with very slow responses (McVay & Kane, 2012b; Unsworth et al., 2010; Unsworth et al., 2012), indicating that individuals with low WM capacity frequently fail to maintain the task goal. In their study, McCabe et al. (2010) examined the relationship between WM capacity and executive functioning using a factor-based analytical approach. They administered several tests to measure WM capacity and executive functioning to more than 200 participants aged between 18 and 90 years. The results indicated a very strong correlation between WM capacity and executive function constructs, demonstrating that they are closely related. These researchers suggest that there is an underlying common component between WM capacity and executive functioning, which they referred to as "executive attention," following the model of Kane and colleagues (Engle & Kane, 2004; Kane & Engle, 2002; McVay & Kane, 2009). This component is essential for maintaining objectives and resolving interference during complex cognitive tasks. On the other hand, Hester and Garavan (2005) conducted a series of three experiments in which they demonstrated that an increase in WM load affects the ability to apply executive control over the elements maintained in WM. In other words, WM load makes executive control over its contents increasingly difficult, particularly with task switching and inhibitory control.

To explain the link between WM and EFs at the neuronal level, many studies using Positron Emission Tomography (PET) and Functional Magnetic Resonance Imaging (fMRI) have been conducted. The results revealed an interaction between executive processes and WM (e.g., Carpenter et al., 2000). Cognitive control and WM share a common neural substrate (Kane & Engle, 2002); the dorsolateral prefrontal cortex (dlPFC), which is known to be central in cognitive control, is also central in WM tasks. In another study of 525 cognitively normal subjects, Bailey et al. (2016) evaluated the *N-back* task of WM using fMRI to determine whether WM activation was associated with performance on executive tasks. The results revealed that activation during the *N-back* task of WM occurs in regions involved in executive functioning.

1. Concepts of Working Memory, Cognitive Flexibility, and Planning

1.1. Definition

1.1.1. Working Memory

Baddeley and Hitch (1974) defined WM as a cognitive mechanism that allows for the temporary storage and manipulation of information. In 1986, Baddeley argued that this system is specifically designed for holding and manipulating information needed to perform complex cognitive tasks, like comprehension, reasoning, and learning. However, in 2003, he redefined it as a limited cognitive structure that links perception, long-term memory, and action. Later, in 2007, Seron offered a definition that was seen as the result of decades of research, emphasizing that WM includes all the mental processes involved in retaining and manipulating information to achieve a goal.

The definitions above suggest that WM is a system that carries out two closely connected activities, enabling it to serve as an executive function:

- Active maintenance: it refers to the activity that controls the content of the information held in mind and blocks access to irrelevant information related to the task in progress. Thus, working memory is considered to have a role in resisting distraction.
- Information processing: it allows the simultaneous manipulation of information during the performance of different cognitive activities.

WM is the most widely used concept to explain the development of children in complex cognitive tasks. The capacity of this system increases both quantitatively and qualitatively during childhood (Molliere, 2013). In other words, as children grow older, their memory span expands, and they develop new strategies. Additionally, WM impacts cognitive functioning, in this sense consistent findings linked it to reading performance (El-Mir, 2017, 2020, 2022; Naciri & El-Mir, 2019), reading comprehension (Bouayad & El-Mir, 2022), and academic achievement (El-Mir, 2019). Its functioning has also been shown to decline in some neurodevelopmental disorders, such as autism (Guennach & El-Mir, 2019) and specific developmental language disorders (Kriblou & El-Mir, 2021, 2024). Research has shown that WM is one of the memory structures most affected by depression (Dahbi & El-Mir, 2020) and aging (El-Mir, 2021). It has also been proved that WM functioning is affected by emotional state (Bousbaïat & El-Mir, 2021; El-Mir, 2018). Furthermore, cognitive training improves WM capacity in children with autism spectrum disorder (Sedjari & El-Mir, 2021; Sedjari, El-Mir & Souirti, 2023), children with attention-deficit/hyperactivity disorder (Alaoui Belghiti & El-Mir, 2023), and children with dyslexia (Ammour & El-Mir, 2023). The effect of cognitive training on working memory has also been confirmed (El-Mir & Sedjari, 2022). Cognitive training is also related to improvement in working memory performance in people with schizophrenia (El-Haddadi & El-Mir, 2022).

Along these lines, Siegler (2005) suggests that the development of WM with age reflects the acquisition of new strategies and the refinement of existing ones. Furthermore, the different components of this function do not develop at the same pace, and their capacity increases linearly between the ages of 4 and 14 (Gathercole, 1999; Gathercole et al., 2004; Nevo & Breznitz, 2013).

1.1.2. Cognitive flexibility

It is a key component in executive functioning. Known as mental flexibility, task switching, or set switching, this ability allows switching between distinct actions or thoughts depending on the situation and the environment (Armbruster et al., 2012, Geurts et al., 2009; Monsell, 2003) and it is essential for adaptive behavior (Badre & Wagner, 2006). It helps to

update contingencies and alternate between responses that guide behavior (Buss & Lowery, 2020; Van der Linden et al., 2000). This function involves changing tasks, focus, or rules (Monsell, 2003) and allows an individual to effectively shift focus from a prior task, adapt by establishing a new set of responses, and apply this updated approach to the current task (Dajani & Lucina, 2015). Enhanced cognitive flexibility is linked to positive outcomes across all stages of life, including improved reading skills in childhood (de Abreu et al., 2014), greater resilience to stress and adverse life events in adulthood (Genet & Siemer, 2011), increased creativity in adults (Chen et al., 2014), and a better quality of life in older age (Davis et al., 2010).

Eslinger and Grattan (1993) distinguish between two types of flexibility: reactive flexibility and spontaneous flexibility. The first refers to adjusting cognition or behavior based on situational demands. According to Slamecka (1968), there are two forms of reactive flexibility: intradimensional flexibility occurs when relevant aspects of a task remain constant, and extradimensional flexibility is involved when a conceptual change in how a task is approached. While spontaneous flexibility consists of generating diverse and new ideas in response to a single question (Zmigrode et al., 2019), it is used when the context is stable and doesn't require modulation, and it is closely tied to the concept of fluency. As indicated by Getzels and Jackson (1962), it can be categorized into two types: ideational fluency (the ability to produce many ideas) and semantic spontaneous flexibility (the ability to generate diverse and varied ideas). Semantic spontaneous flexibility is often associated with divergent thinking, which focuses on producing numerous, relevant, and varied ideas (Chapey, 1994). Cognitive flexibility abilities start emerging in early childhood and show significant increase between the ages of 7 and 9. By the age of 10, this function is mostly developed (Dick, 2014), but it continues to refine and improve during adolescence and adulthood (Anderson, 2002; Hunter & Sparrow, 2012), reaching its highest level between 21 and 30 years of age (Cepeda et al., 2001).

1.1.3. Planning

Planning is one aspect of executive functioning. It is a complex and dynamic cognitive process that involves the evaluation, formulation, and selection of a sequence of actions and thoughts aimed at achieving a goal (Hill, 2004). It requires the ability to organize and plan a series of steps to reach a specific objective (Anderson et al., 2001; Dennis, 2006), conceptualizing changes from the current situation and viewing the environment objectively, managing oneself in relation to the environment, and developing alternative strategies when necessary (Lezak et al., 2004). In other words, it is the skill that helps break down tasks into manageable steps, understand potential challenges, and predict obstacles that could affect task completion (Downing, 2015). As a form of problem-solving, planning also involves mentally executing goal-directed actions to anticipate and assess their potential outcomes (Kaller et al., 2004). Prior to action, the mental representation of the current situation must be transformed into a desired goal state by generating multiple hypothetical scenarios. In addition to these mechanisms, planning requires cognitive abilities such as recognizing goal attainment, anticipating future events related to execution, and storing representations to guide movement from the initial state to the goal (Carlin et al., 2000). Various studies have shown that planning performance increases between the ages of 3 and 14, eventually reaching adult-level efficiency (e.g., Mahone et al., 2002; Malloy-Diniz et al., 2008; Vuelta et al., 2004).

1.2. Neurobiological Bases

1.2.1. Working Memory

Due to a lack of consensus regarding the brain organization of WM, which, according to Rottschy et al. (2012), is attributed to the diversity of tasks and paradigms designed to measure the different aspects of WM, several meta-analyses have been conducted, highlighting the complex and distributed neural networks underlying working memory. Studies show that spatial information activates the superior posterior cortex, while object-related information engages the inferior temporal cortex. Verbal information, however, activates the left lateral inferior frontal cortex and premotor cortex. The superior frontal cortex is involved in tasks requiring information updating or sequence memorization, whereas the inferior prefrontal gyrus is activated during information manipulation tasks (Wager & Smith, 2003). The meta-analysis by Owen et al. (2005) identified activations in frontal and parietal regions, including the median posterior parietal cortex, the bilateral and median premotor cortex, the bilateral rostral PFC, the bilateral dorsolateral prefrontal, and the bilateral mid-ventrolateral prefrontal (vlPFC). Rottschy et al. (2012) found widespread activations in both hemispheres, particularly in the anterior insula, inferior frontal gyrus, and posterior regions like the supplementary motor cortex, intra-parietal sulcus, parietal lobule, ventral visual cortex, and cerebellar V1 lobule. Subcortical activations were also noted in the basal ganglia and thalamus, which connect to the prefrontal and temporal regions. Finally, Yaple and Arsalidou (2018) observed that WM tasks activate posterior brain regions and the right insular cortex.

1.2.2. Cognitive Flexibility

Recent neuroimaging meta-analyses on cognitive flexibility in typically developing adults have reported that task switching probably emerges from the interaction of a network involving specific regions in the frontal and parietal cortices (e.g., Kim et al., 2012; Leber et al., 2008; Niendam et al., 2012; Schmitz et al., 2006; Zühlsdorff et al., 2023). This network includes complex cortical regions such as the vlPFC, dlPFC, anterior cingulate cortex, right anterior insula, as well as the premotor cortex, inferior and superior parietal cortices, inferior temporal cortex, occipital cortex, and subcortical structures like the caudate and thalamus. Additionally, these changes are accompanied by reduced connectivity between the anterior insula, orbitofrontal cortex, and occipital cortex (Zühlsdorff et al., 2023).

1.2.3. Planning

Findings from lesion studies, pathological research, and neuroimaging studies using the *Tower of London (TOL)* task strongly highlight the critical role of the prefrontal cortex (PFC), particularly its dorsolateral and rostral regions, in complex problem-solving (Albert & Steinberg, 2011). Using this task, activation was detected across a widespread network of cortical regions, including the prefrontal, cingulate, premotor, parietal, and occipital cortices (Baker et al., 1996). Dagher et al. (1999) indicated that the neural regions involved in planning include the prefrontal cortex, particularly the dlPFC, as well as the anterior cingulate cortex, parietal cortex, and caudate nucleus, which are critical for coordinating complex cognitive processes such as goal setting, strategy formation, and evaluating potential outcomes during planning tasks. As for Van den Heuvel et al. (2003), planning activity was associated with activation in the frontostriatal, visuospatial, and motor systems, specifically involving dlPFC, anterior prefrontal cortex, striatum, precuneus, inferior parietal cortex, premotor cortex, and the Supplementary motor area.

2. Working Memory and Cognitive Flexibility

In Baddeley's (1986) multi-component model, one of the main functions of the central executive is the ability to switch efficiently between tasks (Vandierendonck, 2016). According to Diamond (2013), cognitive flexibility consists of updating information in WM to identify the most appropriate response to the current situation. Several theories have pointed to a strong link between WM and cognitive flexibility (Mayr & Kliegl, 2003; Meiran & Kessler, 2008; Sohn & Anderson, 2001). More specifically, research has shown that this function is closely related to verbal WM, particularly the phonological loop in Baddeley's model. On one hand, it has been shown that verbalizing task goals can improve cognitive flexibility (e.g., Goshke, 2000). On the other hand, depletion of verbal WM can lead to errors and slower processing (Saeki & Saito, 2009; Saeki et al., 2006), affecting the cost of this function (e.g., Baddeley et al., 2001; Miyake et al., 2004; Saeki & Saito, 2004). For example, Baddeley et al. (2001) observed that the cost associated with task switching increased when it was combined with a verbal task. A study by Souza et al. (2012) investigated how WM load impacts task switching. Participants were asked to memorize one to three sets of numbers and complete several tasks, including determining whether the number was greater or less than five, whether it was odd or even, or its position on a number line (inside or outside a range). The results revealed that as the WM load increased, so did the cost of switching tasks, particularly when participants were asked to remember three lists instead of just one or two. This indicates that when WM load is higher, there is a greater need for articulatory rehearsal, which increases the task switching cost, as suggested in earlier studies (Baddeley et al., 2001). Likewise, other studies have shown that increased task switching demands can impair WM performance (Liefoghe et al., 2008). Butler et al. (2011) found a notable link between WM capacity and task switching costs, with higher task-switching costs associated with lower WM capacity. However, some studies have not found a similar relationship, suggesting that WM capacity may not always be closely linked with cognitive flexibility (Draheim et al., 2016; Hambriek & Altmann, 2015; Kane et al., 2007; Logan, 2004; Pettigrew & Martim, 2016; Unsworth & Engle, 2008). Some researchers even argue that WM should be considered a distinct construct from other executive functions, with no clear connection to task switching (Logan, 2004; Oberauer et al., 2007; Oberauer, 2009; Vandierendonck, 2016).

At the neuronal level, a meta-analysis of 193 neuroimaging studies based on the use of different measures of executive functioning (including WM and task switching) revealed that a network of frontal and parietal regions was constantly active in all areas of the EFs examined. This network included the cingulate cortex anterior, the dlPFC in the frontal lobes, and inferior and superior parietal lobes (Niendam et al., 2012). Despite the differences in modalities, many studies have shown that WM tasks systematically activate the frontoparietal network called the cognitive control network (Kondo et al., 2004; Osaka et al., 2003; Osaka et al., 2004; Owen et al., 2005; Rottschy et al., 2012; Van der Linden et al., 2007; Wager & Smith, 2003), and the performance of task switching is also associated with the activation of this network (Dajani & Uddin, 2015; Richter & Yeung, 2014; Ruge et al., 2013). Other studies on cognitive flexibility reported activation in dlPFC and vlPFC, the additional motor area, lower cingulate cortex, and upper and lower parietal cortex (Karayanidis et al., 2010). These regions are similar to those reported in the WM neuroimaging literature.

3. Working Memory and Planning

In their founding book, *Plans and the Structure of Behavior* (Miller et al., 1960), WM is used for the execution of plans. Nevertheless, many authors have linked WM and planning. It has been shown that people with high WM ability are significantly better at tasks traditionally used to evaluate planning (e.g. Gihooley et al., 2002; Miyake et al., 2001; Zook et

al., 2004). Ohbayashi et al. (2003) reported that during movement planning, WM plays a crucial role in actively maintaining relevant information and then converting it into a movement program to achieve goals. Another study by Spiegel et al. (2012) showed that the planning phase and WM share cognitive resources, and both functions seem fundamental to organizing action. Altagassen et al. (2007) revealed that in healthy people, each of the three components of WM was linked to the task of the *TOL*. Studies have indicated that the phonological loop helps to achieve verbally generated mental plans (Altagassen et al., 2007), the visuospatial sketchpad is important in the construction and reformulation of plans (Phillips, 1999), while the central executive may be involved in monitoring the achievement of objectives and the change of focus. Episodic buffer, and despite the absence of studies on its role in planning tasks (Altagassen et al., 2007), it appears to play an important role as the *ToL* task requires the development of strategic information in long-term memory and its subsequent retrieval.

In 2014, Behmer and Fournier conducted a study of 168 participants to examine whether the ability to plan an action and maintain a plan can be influenced by the WM capacity. Participants were divided into two groups: participants with low WM capacity and others with high WM capacity. All participants were asked to briefly maintain a stimulus action plan in WM that they would execute immediately after responding to another stimulus. First, arrows pointing left or right are shown on the screen with an asterisk above or below the point of the arrow. The arrow points indicate which hand the participant should use (left or right), and the asterisk indicates the direction of movement (upper or lower key). There were four different action plans (left hand moved up, left hand moved down, right hand moved up, and right hand moved down). Participants should use the correct keyboard buttons to indicate the correct answer (the correct plan), but before answering this part of the experiment they should double-press the left key on the keyboard with their left hand if they see a green numeric symbol (#) and double-press the right key with their right hand if the symbol is red. The results of this study indicated that participants with low WM capacity did not remember the action plan from the first part of the experiment, as well as those with high WM capacity.

Anatomically, many researchers have conducted studies using fMRI, PET and mono-photon emission tomography to specify brain areas associated with planning ability (e.g., Baker et al., 1996; Dagher et al., 1999; Morris et al., 1993; Rowe et al., 2001). Although it is difficult to determine the areas involved in each cognitive process (Unterrainer & Owen, 2006), these researchers indicated that the rostralateral and dlPFC, premotor cortex, parietal cortex, and inferior frontal gyrus are among the active zones associated with the planning task (*ToL*). These areas were also linked to WM tasks (Owen et al., 2005; Rottschy et al., 2012; Wager & Smith, 2003). Again, in a recent study, Pouladi et al. (2021) put 15 subjects under transcranial direct current stimulation to assess whether dlPFC stimulation will lead to improved WM and planning. Using the *N-back* task and the *ToL* task, these researchers showed that both functions improve after stimulation of this region, which proves that WM and planning share a common neural substrate called the dlPFC.

Discussion

The present article specifically focuses on the neurocognitive relationship between WM and two key EFs: cognitive flexibility and planning. Indeed, the relationship between WM and EFs is well-established (McCabe et al., 2010), despite the heterogeneity of results observed in the literature. This variability can be attributed to multiple factors, such as differences in participant characteristics, methodological approaches used to measure these constructs, and the inherent complexity of executive functioning. WM is considered a core component of cognitive control and a strong predictor of high-level cognition. Individuals with high WM capacity tend to achieve higher scores in complex cognitive tasks (e.g., Redick

et al., 2012). Also, distinct EFs and WM are closely interrelated, sharing underlying neural substrates and activating overlapping brain regions (e.g., Anderson et al., 2001; Huizinga et al., 2006; Lehto et al., 2003; Miyake et al., 2000).

The findings regarding the close relationship between WM and other EFs can serve as a foundation for improving the efficiency of neurocognitive interventions. For instance, rather than attempting cognitive remediation for multiple impaired EFs simultaneously, it may be more effective to focus on remediating WM alone. Given the central role of its capacity in cognitive functioning, improvement of WM may have an impact on other EFs, enhancing the efficiency of interventions (Sedjari & El-Mir, 2021, Sedjari et al., 2023).

Conclusion

In sum, WM, cognitive flexibility, and planning are distinct but interrelated executive components that share underlying neural substrates, generating activation in overlapping brain regions. This conclusion supports the idea that WM could be a prime target for neurocognitive therapies, allowing an efficient and practical means to enhance not just its capacity but also other EFs that rely on it.

References

- Ackerman, P. L., Beier, M. E., & Boyle, M. O. (2005). Working memory and intelligence: The same or different constructs? *Psychological bulletin*, *131*(1), 30. <https://doi.org/10.1037/0033-2909.131.1.30>
- Albert, D., & Steinberg, L. (2011). Age Differences in Strategic Planning as Indexed by the Tower of London. *Child Development*, *82*(5), 1501-1517. <https://doi.org/10.1111/j.1467-8624.2011.01613.x>
- Altgassen, M., Phillips, L., Kopp, U., & Kliegel, M. (2007). Role of working memory components in planning performance of individuals with Parkinson's disease. *Neuropsychologia*, *45*(10), 2393-2397. <https://doi.org/10.1016/j.neuropsychologia.2007.02.018>
- Anderson, P. (2002). Assessment and Development of Executive Function (EF) During Childhood. *Child Neuropsychology*, *8*(2), 71-82. <https://doi.org/10.1076/chin.8.2.71.8724>
- Armbruster, D. J. N., Ueltzhöffer, K., Basten, U., & Fiebach, C. J. (2012). Prefrontal Cortical Mechanisms Underlying Individual Differences in Cognitive Flexibility and Stability. *Journal Of Cognitive Neuroscience*, *24*(12), 2385-2399. https://doi.org/10.1162/jocn_a_00286
- Baddeley, A. D. (1986). *Working Memory*. New York: Oxford University Press. <https://studylib.net/doc/7657654/baddeley--a.--1986--working-memory.-oxford--oxford-unive>.
- Baddeley, A., Chincotta, D., & Adlam, A. (2001). Working memory and the control of action: evidence from task switching. *Journal of experimental psychology: General*, *130*(4), 641. <https://doi.org/10.1037/0096-3445.130.4.641>
- Bailey, K., Mlynarczyk, G., & West, R. (2016). Slow wave activity related to working memory maintenance in the N-back task. *Journal of Psychophysiology*. *30*(4), 141-154. <https://doi.org/10.1027/0269-8803/a000164>
- Baker, S. C., Rogers, R. D., Owen, A. M., Frith, C. D., Dolan, R. J., Frackowiak, R. S. J., & Robbins, T. W. (1996). Neural systems engaged by planning: a PET study of the Tower of London task. *Neuropsychologia*, *34*(6), 515-526. [https://doi.org/10.1016/0028-3932\(95\)00133-6](https://doi.org/10.1016/0028-3932(95)00133-6)
- Badre, D., & Wagner, A. D. (2006). Computational and neurobiological mechanisms underlying cognitive flexibility. *Proceedings Of The National Academy Of Sciences*, *103*(18), 7186-7191. <https://doi.org/10.1073/pnas.0509550103>
- Behmer, L. P., Jr, & Fournier, L. R. (2014). Working memory modulates neural efficiency over motor components during a novel action planning task: an EEG study. *Behavioural brain research*, *260*, 1-7. <https://doi.org/10.1016/j.bbr.2013.11.031>
- Bouayad, M. & El-Mir, M. (2022). The impact of executive functions on reading comprehension. *Arab Journal of Psychology*, *7*(1), 125-144. <https://doi.org/10.6084/m9.figshare.21084871.v1>
- Bousbaïat, O. & El-Mir, M. (2021). The Effect of Fear on Working Memory in children. *Arab Journal of Psychology*, *6*(1), 125-138. <https://doi.org/10.6084/m9.figshare.21151573.v1>
- Butler, K. M., Arrington, C. M., & Weywadt, C. (2011). Working memory capacity modulates task performance but has little influence on task choice. *Memory & Cognition*, *39*, 708-724. <https://doi.org/10.3758/s13421-010-0055-y>
- Carpenter, P. A., Just, M. A., & Reichle, E. D. (2000). Working memory and executive function: Evidence from neuroimaging. *Current opinion in neurobiology*, *10*(2), 195-199. [https://doi.org/10.1016/S0959-4388\(00\)00074-X](https://doi.org/10.1016/S0959-4388(00)00074-X)
- Cepeda, N. J., Kramer, A. F., & Gonzalez de Sather, J. C. M. (2001). Changes in executive control across the life span: Examination of task-switching performance. *Developmental Psychology*, *37*(5), 715-730. <https://doi.org/10.1037/0012-1649.37.5.715>
- Chen, Q., Yang, W., Li, W., Wei, D., Li, H., Lei, Q., Zhang, Q., & Qiu, J. (2014). Association of creative achievement with cognitive flexibility by a combined voxel-based morphometry and resting-state functional connectivity study. *NeuroImage*, *102*, 474-483. <https://doi.org/10.1016/j.neuroimage.2014.08.008>
- Conway, A. R., Cowan, N., & Bunting, M. F. (2001). The cocktail party phenomenon revisited: The importance of working memory capacity. *Psychonomic bulletin & review*, *8*, 331-335. <https://doi.org/10.3758/BF03196169>

- Conway, A. R., Kane, M. J., Bunting, M. F., Hambrick, D. Z., Wilhelm, O., & Engle, R. W. (2005). Working memory span tasks: A methodological review and user's guide. *Psychonomic bulletin & review*, *12*, 769-786. <https://doi.org/10.3758/BF03196772>
- Cowan, N. (1995). *Verbal Working Memory: A View with a Room*. University of Illinois Press. <https://doi.org/10.2307/1423105>
- Dagher, A., Owen, A. M., Boecker, H., & Brooks, D. J. (1999). Mapping the network for planning: a correlational PET activation study with the Tower of London task. *Brain*, *122*(10), 1973-1987. <https://doi.org/10.1093/brain/122.10.1973>
- Dahbi S. & El-Mir, M. (2020). Impact of depression on working memory: Comparative study of the functioning of working memory in a group of patients with characterized depressive disorder and a control group. *Arab Journal of Psychology*, *5*(2), 178-188. <https://doi.org/10.6084/m9.figshare.21151630.v1>
- Dajani, D. R., & Uddin, L. Q. (2015). Demystifying cognitive flexibility: Implications for clinical and developmental neuroscience. *Trends in neurosciences*, *38*(9), 571-578. <https://doi.org/10.1016/j.tins.2015.07.003>
- De Abreu, P. M. J. E., Abreu, N., Nikaedo, C. C., Puglisi, M. L., Tourinho, C. J., Miranda, M. C., Befi-Lopes, D. M., Bueno, O. F. A., & Martin, R. (2014). Executive functioning and reading achievement in school: a study of Brazilian children assessed by their teachers as "poor readers". *Frontiers In Psychology*, *5*. <https://doi.org/10.3389/fpsyg.2014.00550>
- Diamond, A. (2013). Executive functions. *Annual review of psychology*, *64*, 135-168. <https://doi.org/10.1146/annurev-psych-113011-143750>
- Draheim, C., Hicks, K. L., & Engle, R. W. (2016). Combining reaction time and accuracy: The relationship between working memory capacity and task switching as a case example. *Perspectives on Psychological Science*, *11*(1), 133-155. <https://doi.org/10.1177/1745691615596990>
- Davis, S., Rawlings, B., Clegg, J. M., Ikejimba, D., Watson-Jones, R. E., Whiten, A., & Legare, C. H. (2022). Cognitive flexibility supports the development of cumulative cultural learning in children. *Scientific Reports*, *12*(1). <https://doi.org/10.1038/s41598-022-18231-7>
- Dick, A. S. (2014). The development of cognitive flexibility beyond the preschool period : An investigation using a modified Flexible Item Selection Task. *Journal Of Experimental Child Psychology*, *125*, 13-34. <https://doi.org/10.1016/j.jecp.2014.01.021>
- El-Haddadi, A. & El-Mir, M. (2022). Working memory training in patients with schizophrenia. *Arab Journal of Psychology*, *7*(1). 41-53. <https://doi.org/10.6084/m9.figshare.21084868.v2>
- El-Mir, M. (2017). The effect of working memory capacity on word recognition speed in Arabic second grade readers. *Arab Journal of Psychology*, *3*(1), 149-160. <https://doi.org/10.6084/m9.figshare.12155970.v1>
- El-Mir, M. (2018). The role of emotional processes in memory functioning. *Arab Journal of Psychology*, *3*(2), 94-103. <https://doi.org/10.6084/m9.figshare.12155955.v1>
- El-Mir, M. (2019). Impact of memory on school performance. *Arab Journal of Psychology*, *4*(2), 184-196. <https://doi.org/10.6084/m9.figshare.12152199.v1>
- El-Mir, M. (2020). Effect of working memory capacity on Arabic reading development in primary school pupils in Morocco. *Arab Journal of Psychology*, *5*(1), 92-106. <https://doi.org/10.6084/m9.figshare.21586932.v1>
- El-Mir, M. (2021). Memory functioning in aging. *Nafssaniat*, *73*, 17-29. <https://doi.org/10.6084/m9.figshare.21151552.v1>
- El-Mir, M. (2022). *Reading and working memory*. Books Cultural Center: Casablanca, Beirut, ISBN: 978-9920-677-25-7. <https://doi.org/10.6084/m9.figshare.21096664.v1>
- El-Mir, M. & Sedjari, S. (2022). Effect of working memory training on mental disorders. *Arab Journal of Psychology*, *7*(3). 91-106. <https://doi.org/10.57642/AJOPSY8>
- Guennach, A. & El-Mir, M. (2019). Autism spectrum disorder and working memory: A comparative study between children with ASD and normal children. *Arab Journal of Psychology*, *4*(2), 123-133. <https://doi.org/10.6084/m9.figshare.12155694.v2>
- Engle, R. W., & Kane, M. J. (2004). Executive Attention, Working Memory Capacity, and a Two-Factor Theory of Cognitive Control. In B. H. Ross (Ed.), *The psychology of learning and*

- motivation: Advances in research and theory* (pp. 145–199). Elsevier Science. <https://psycnet.apa.org/record/2004-00153-005>
- Genet, J. J., & Siemer, M. (2010). Flexible control in processing affective and non-affective material predicts individual differences in trait resilience. *Cognition & Emotion*, 25(2), 380-388. <https://doi.org/10.1080/02699931.2010.491647>
- Ger, E., & Roebers, C. M. (2023). The Relationship between Executive Functions, Working Memory, and Intelligence in Kindergarten Children. *Journal of Intelligence*, 11(4), 64. <https://doi.org/10.3390/jintelligence11040064>
- Gilhooly, K. J., Wynn, V., Phillips, L. H., Logie, R. H., & Sala, S. D. (2002). Visuo-spatial and verbal working memory in the five-disc Tower of London task: An individual differences approach. *Thinking & reasoning*, 8(3), 165-178. <https://doi.org/10.1080/13546780244000006>
- Goschke, T. (2000). Reconfiguration of stimulus task sets and response task sets during task switching. *Control of cognitive processes*, 331. https://www.researchgate.net/publication/38140172_Intentional_reconfiguration_and_involuntary_persistence_in_task-set_switching
- Hambrick, D. Z., & Altmann, E. M. (2015). The role of placekeeping ability in fluid intelligence. *Psychonomic bulletin & Review*, 22, 1104-1110. <https://doi.org/10.3758/s13423-014-0764-5>
- Hunter, S. J., & Sparrow, E. P. (Eds.). (2012). *Executive function and dysfunction: Identification, assessment and treatment*. Cambridge University Press.
- Hester, R., & Garavan, H. (2005). Working memory and executive function: The influence of content and load on the control of attention. *Memory & cognition*, 33, 221-233. <https://doi.org/10.3758/BF03195311>
- Kane, M. J., Conway, A. R. A., Hambrick, D. Z., & Engle, R. W. (2007). Variation in working memory capacity as variation in executive attention and control. In A. R. A. Conway, C. Jarrold, M. J. Kane (Eds.), *Variation in working memory* (pp. 21-46). Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780195168648.003.0002>
- Kane, M. J., & Engle, R. W. (2002). The role of prefrontal cortex in working-memory capacity, executive attention, and general fluid intelligence: An individual-differences perspective. *Psychonomic bulletin & review*, 9(4), 637-671. <https://doi.org/10.3758/BF03196323>
- Karayanidis, F., Jamadar, S., Ruge, H., Phillips, N., Heathcote, A., & Forstmann, B. U. (2010). Advance preparation in task-switching: converging evidence from behavioral, brain activation, and model-based approaches. *Frontiers in Psychology*, 1, 25. <https://doi.org/10.3389/fpsyg.2010.00025>
- Kim, P., Jenkins, S. E., Connolly, M. E., Deveney, C. M., Fromm, S. J., Brotman, M. A., Nelson, E. E., Pine, D. S., & Leibenluft, E. (2011). Neural correlates of cognitive flexibility in children at risk for bipolar disorder. *Journal Of Psychiatric Research*, 46(1), 22-30. <https://doi.org/10.1016/j.jpsychires.2011.09.015>
- Kimberg, D. Y., D'esposito, M., & Farah, M. J. (1997). Cognitive functions in the prefrontal cortex—Working memory and executive control. *Current Directions in Psychological Science*, 6(6), 185-192. <https://doi.org/10.1111/1467-8721.ep10772959>
- Kondo, H., Morishita, M., Osaka, N., Osaka, M., Fukuyama, H., & Shibasaki, H. (2004). Functional roles of the cingulo-frontal network in performance on working memory. *Neuroimage*, 21(1), 2-14. <https://doi.org/10.1016/j.neuroimage.2003.09.046>
- Kriblou A. & El-Mir, M. (2021). Working memory functioning in children with specific developmental language disorders. *Arab Journal of Psychology*, 6(2), 17-32. <https://doi.org/10.6084/m9.figshare.21151540.v1>
- Kriblou A. & El-Mir, M. (2024). The Effect of Cognitive Training on Working Memory Performance in Verbal Tasks in Children with Specific Developmental Language Disorders. *Arab Journal of Psychology*, 9(2), 6-17. <https://doi.org/10.57642/AJOPSY913>
- Leber, A. B., Turk-Browne, N. B., & Chun, M. M. (2008). Neural predictors of moment-to-moment fluctuations in cognitive flexibility. *Proceedings Of The National Academy Of Sciences*, 105(36), 13592-13597. <https://doi.org/10.1073/pnas.0805423105>
- Liefooghe, B., Barrouillet, P., Vandierendonck, A., & Camos, V. (2008). Working memory costs of task switching. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 34(3), 478–494. <https://doi.org/10.1037/0278-7393.34.3.478>

- Logan, G. D. (2004). Working Memory, Task Switching, and Executive Control in the Task Span Procedure. *Journal of Experimental Psychology: General*, 133(2), 218–236. <https://doi.org/10.1037/0096-3445.133.2.218>
- Mayr, U., & Kliegl, R. (2003). Differential effects of cue changes and task changes on task-set selection costs. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 29(3), 362–372. <https://doi.org/10.1037/0278-7393.29.3.362>
- McCabe, D. P., Roediger, H. L. III, McDaniel, M. A., Balota, D. A., & Hambrick, D. Z. (2010). The relationship between working memory capacity and executive functioning: Evidence for a common executive attention construct. *Neuropsychology*, 24(2), 222–243. <https://doi.org/10.1037/a0017619>
- McVay, J. C., & Kane, M. J. (2009). Conducting the train of thought: Working memory capacity, goal neglect, and mind wandering in an executive-control task. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 35(1), 196–204. <https://doi.org/10.1037/a0014104>
- McVay, J. C., & Kane, M. J. (2012a). Drifting from slow to “d'oh!": Working memory capacity and mind wandering predict extreme reaction times and executive control errors. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 38(3), 525–549. <https://doi.org/10.1037/a0025896>
- McVay, J. C., & Kane, M. J. (2012b). Why does working memory capacity predict variation in reading comprehension? On the influence of mind wandering and executive attention. *Journal of Experimental Psychology: General*, 141(2), 302–320. <https://doi.org/10.1037/a0025250>
- Meiran, N., & Kessler, Y. (2008). The task rule congruency effect in task switching reflects activated long-term memory. *Journal of Experimental Psychology: Human Perception and Performance*, 34(1), 137–157. <https://doi.org/10.1037/0096-1523.34.1.137>
- Miller, G. A., Galanter, E., & Pribram, K. H. (1960). *Plans and the structure of behavior*. Henry Holt and Co. <https://doi.org/10.1037/10039-000>
- Miyake, A., Emerson, M. J., Padilla, F., & Ahn, J. C. (2004). Inner speech as a retrieval aid for task goals: The effects of cue type and articulatory suppression in the random task cuing paradigm. *Acta psychologica*, 115(2-3), 123–142. <https://doi.org/10.1016/j.actpsy.2003.12.004>
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. *Cognitive psychology*, 41(1), 49–100. <https://doi.org/10.1006/cogp.1999.0734>
- Miyake, A., Friedman, N. P., Rettinger, D. A., Shah, P., & Hegarty, M. (2001). How are visuospatial working memory, executive functioning, and spatial abilities related? A latent-variable analysis. *Journal of Experimental Psychology: General*, 130(4), 621–640. <https://doi.org/10.1037/0096-3445.130.4.621>
- Morris, R. G., Ahmed, S., Syed, G. M., & Toone, B. K. (1993). Neural correlates of planning ability: frontal lobe activation during the Tower of London test. *Neuropsychologia*, 31(12), 1367–1378. [https://doi.org/10.1016/0028-3932\(93\)90104-8](https://doi.org/10.1016/0028-3932(93)90104-8)
- Naciri, M. & El-Mir, M. (2019). Reading and phonologico-morphological characteristics of Arabic: a comparative study of good readers and dyslexics. *Arab Journal of Psychology*, 4(1), 67–79. <https://doi.org/10.6084/m9.figshare.12155937.v1>
- Norman, D. A., & Shallice, T. (1986). *Attention to action: Willed and automatic control of behavior*. Springer US. https://doi.org/10.1007/978-1-4757-0629-1_1
- Niendam, T. A., Laird, A. R., Ray, K. L., Dean, Y. M., Glahn, D. C., & Carter, C. S. (2012). Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. *Cognitive, Affective, & Behavioral Neuroscience*, 12, 241–268. <https://doi.org/10.3758/s13415-011-0083-5>
- Oberauer, K. (2009). Design for a working memory. *Psychology of learning and motivation*, 51, 45–100. [https://doi.org/10.1016/S0079-7421\(09\)51002-X](https://doi.org/10.1016/S0079-7421(09)51002-X)
- Oberauer, K., Süß, H.-M., Wilhelm, O., & Sander, N. (2007). Individual differences in working memory capacity and reasoning ability. In A. R. A. Conway, C. Jarrold, M. J. Kane (Eds.) & A. Miyake & J. N. Towse (Ed.), *Variation in working memory* (pp. 49–75). Oxford University Press. <http://dx.doi.org/10.1093/acprof:oso/9780195168648.003.0003>

- Ohbayashi, M., Ohki, K., & Miyashita, Y. (2003). Conversion of working memory to motor sequence in the monkey premotor cortex. *Science*, *301*(5630), 233-236. <https://doi.org/10.1126/science.1084884>
- Osaka, M., Osaka, N., Kondo, H., Morishita, M., Fukuyama, H., Aso, T., & Shibasaki, H. (2003). The neural basis of individual differences in working memory capacity: an fMRI study. *NeuroImage*, *18*(3), 789-797. [https://doi.org/10.1016/S1053-8119\(02\)00032-0](https://doi.org/10.1016/S1053-8119(02)00032-0)
- Osaka, N., Osaka, M., Kondo, H., Morishita, M., Fukuyama, H., & Shibasaki, H. (2004). The neural basis of executive function in working memory: an fMRI study based on individual differences. *Neuroimage*, *21*(2), 623-631. <https://doi.org/10.1016/j.neuroimage.2003.09.069>
- Owen, A. M., McMillan, K. M., Laird, A. R., & Bullmore, E. (2005). N-back working memory paradigm: A meta-analysis of normative functional neuroimaging studies. *Human brain mapping*, *25*(1), 46-59. <https://doi.org/10.1002/hbm.20131>
- Pettigrew, C., & Martin, R. C. (2016). The role of working memory capacity and interference resolution mechanisms in task switching. *Quarterly Journal of Experimental Psychology*, *69*(12), 2431-2451. <https://doi.org/10.1080/17470218.2015.1121282>
- Phillips, L. H. (1999). The role of memory in the Tower of London task. *Memory*, *7*(2), 209-231. <https://doi.org/10.1080/741944066>
- Pouladi, F., Bagheri, M., Askarizadeh, G., & Moradi, A. (2021). Stimulation of the dorsolateral-prefrontal cortex improves working memory and planning. *Cognition, Brain, Behavior*, *25*(1), 1-17. <https://doi.org/10.24193/cbb.2021.25.01>
- Redick, T. S., Calvo, A., Gay, C. E., & Engle, R. W. (2011). Working memory capacity and go/no-go task performance: Selective effects of updating, maintenance, and inhibition. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *37*(2), 308-324. <https://doi.org/10.1037/a0022216>
- Redick, T. S., Unsworth, N., Kelly, A. J., & Engle, R. W. (2012). Faster, smarter? Working memory capacity and perceptual speed in relation to fluid intelligence. *Journal of Cognitive Psychology*, *24*(7), 844-854. <https://doi.org/10.1080/20445911.2012.704359>
- Richter, F. R., & Yeung, N. (2014). Neuroimaging studies of task switching. In J. A. Grange & G. Houghton (Eds.), *Task switching and cognitive control* (pp. 237-271). Oxford University Press. <https://doi.org/10.1093/acprof:osobl/9780199921959.003.0010>
- Rottschy, C., Langner, R., Dogan, I., Reetz, K., Laird, A. R., Schulz, J. B., Fox, P. T., & Eickhoff, S. B. (2012). Modelling neural correlates of working memory: a coordinate-based meta-analysis. *NeuroImage*, *60*(1), 830-846. <https://doi.org/10.1016/j.neuroimage.2011.11.050>
- Rowe, J. B., Owen, A. M., Johnsrude, I. S., & Passingham, R. E. (2001). Imaging the mental components of a planning task. *Neuropsychologia*, *39*(3), 315-327. [https://doi.org/10.1016/S0028-3932\(00\)00109-3](https://doi.org/10.1016/S0028-3932(00)00109-3)
- Ruge, H., Jamadar, S., Zimmermann, U., & Karayanidis, F. (2013). The many faces of preparatory control in task switching: reviewing a decade of fMRI research. *Human brain mapping*, *34*(1), 12-35. <https://doi.org/10.1002/hbm.21420>
- Saeki, E., & Saito, S. (2004). Effect of articulatory suppression on task-switching performance: Implications for models of working memory. *Memory*, *12*(3), 257-271. <https://doi.org/10.1080/09658210244000649>
- Saeki, E., & Saito, S. (2009). Verbal representation in task order control: An examination with transition and task cues in random task switching. *Memory & Cognition*, *37*(7), 1040-1050. <https://doi.org/10.3758/MC.37.7.1040>
- Saeki, E., Saito, S., & Kawaguchi, J. (2006). Effects of response-stimulus interval manipulation and articulatory suppression on task switching. *Memory*, *14*(8), 965-976. <https://doi.org/10.1080/09658210601008973>
- Schelble, J. L., Theriault, D. J., & Miller, M. D. (2012). Classifying retrieval strategies as a function of working memory. *Memory & cognition*, *40*, 218-230. <https://doi.org/10.3758/s13421-011-0149-1>
- Schmitz, N., Rubia, K., Daly, E., Smith, A., Williams, S., & Murphy, D. G. (2005). Neural Correlates of Executive Function in Autistic Spectrum Disorders. *Biological Psychiatry*, *59*(1), 7-16. <https://doi.org/10.1016/j.biopsych.2005.06.007>

- Sedjari, S. & El-Mir, M. (2021). Working memory training in autism spectrum disorder. *Arab Journal of Psychology*, 6(1), 194-209. <https://doi.org/10.6084/m9.figshare.21151609.v1>
- Sedjari, S., El-Mir, M., & Souirti, Z. (2023). Working memory training in autism: Near and far transfer. *PubMed*, 101(12), 884-890. <https://pubmed.ncbi.nlm.nih.gov/38477195>
- Sohn, M.-H., & Anderson, J. R. (2001). Task preparation and task repetition: Two-component model of task switching. *Journal of Experimental Psychology: General*, 130(4), 764-778. <https://doi.org/10.1037/0096-3445.130.4.764>
- Souza, A. S., Oberauer, K., Gade, M., & Druery, M. D. (2012). Processing of representations in declarative and procedural working memory. *Quarterly Journal of Experimental Psychology*, 65(5), 1006-1033. <https://doi.org/10.1080/17470218.2011.640403>
- Spiegel, M. A., Koester, D., Weigelt, M., & Schack, T. (2012). The costs of changing an intended action: Movement planning, but not execution, interferes with verbal working memory. *Neuroscience Letters*, 509(2), 82-86. <https://doi.org/10.1016/j.neulet.2011.12.033>
- Unsworth, N., & Engle, R. W. (2008). Speed and accuracy of accessing information in working memory: An individual differences investigation of focus switching. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 34(3), 616-630. <https://doi.org/10.1037/0278-7393.34.3.616>
- Unsworth, N., Heitz, R. P., & Engle, R. W. (2005). Working Memory Capacity in Hot and Cold Cognition. In R. W. Engle, G. Sedek, U. von Hecker, & D. N. McIntosh (Eds.), *Cognitive limitations in aging and psychopathology* (pp. 19-43). Cambridge University Press. <https://doi.org/10.1017/CBO9780511720413.003>
- Unsworth, N., Redick, T. S., Spillers, G. J., & Brewer, G. A. (2012). Variation in working memory capacity and cognitive control: Goal maintenance and microadjustments of control. *Quarterly Journal of Experimental Psychology*, 65(2), 326-355. <https://doi.org/10.1080/17470218.2011.597865>
- Unsworth, N., Spillers, G. J., & Brewer, G. A. (2010). The contributions of primary and secondary memory to working memory capacity: An individual differences analysis of immediate free recall. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 36(1), 240-247. <https://doi.org/10.1037/a0017739>
- Unterrainer, J. M., & Owen, A. M. (2006). Planning and problem solving: from neuropsychology to functional neuroimaging. *Journal of Physiology-Paris*, 99(4-6), 308-317. <https://doi.org/10.1016/j.jphysparis.2006.03.014>
- Van Den Heuvel, O. A., Groenewegen, H. J., Barkhof, F., Lazeron, R. H., Van Dyck, R., & Veltman, D. J. (2003). Frontostriatal system in planning complexity : a parametric functional magnetic resonance version of tower of london task. *NeuroImage*, 18(2), 367-374. [https://doi.org/10.1016/s1053-8119\(02\)00010-1](https://doi.org/10.1016/s1053-8119(02)00010-1)
- van der Linden, W. J. (2007). A hierarchical framework for modeling speed and accuracy on test items. *Psychometrika*, 72(3), 287-308. <https://doi.org/10.1007/s11336-006-1478-z>
- Vandierendonck, A. (2016). A working memory system with distributed executive control. *Perspectives on Psychological Science*, 11(1), 74-100. <https://doi.org/10.1177/1745691615596790>
- Wager, T. D., & Smith, E. E. (2003). Neuroimaging studies of working memory. *Cognitive, Affective, & Behavioral Neuroscience*, 3(4), 255-274. <https://doi.org/10.3758/CABN.3.4.255>
- Wilhelm, O., Hildebrandt, A., & Oberauer, K. (2013). What is working memory capacity, and how can we measure it?. *Frontiers in psychology*, 4, 433. <https://doi.org/10.3389/fpsyg.2013.00433>
- Zook, N. A., Davalos, D. B., DeLosh, E. L., & Davis, H. P. (2004). Working memory, inhibition, and fluid intelligence as predictors of performance on Tower of Hanoi and London tasks. *Brain and cognition*, 56(3), 286-292. <https://doi.org/10.1016/j.bandc.2004.07.003>
- Zühlsdorff, K., Dalley, J. W., Robbins, T. W., & Morein-Zamir, S. (2022). Cognitive flexibility: neurobehavioral correlates of changing one's mind. *Cerebral Cortex*, 33(9), 5436-5446. <https://doi.org/10.1093/cercor/bhac431>